

CONSTRUCTION OF A RECOMBINANT BACMID DNA ENCODING VIRAL PROTEIN-2 OF CANINE PARVOVIRUS USING SITE-SPECIFIC TRANSPOSITION MECHANISM

Mohammad Sadegh Hashemzadeh¹, Seyed Jafar Mousavy^{1*}, Ruhollah Dorostkar², Fatemeh Fotouhi³, Firouz Ebrahimi¹

¹Department of Biology, Faculty of Basic Sciences, Imam Hossein University, Tehran, IRAN

²Applied Virology Research Center, Baqiyatallah University of Medical Sciences, Tehran, IRAN

³Department of Influenza and other Respiratory viruses, Pasteur Institute of Iran, Tehran, IRAN

ABSTRACT

Background and Aim: Canine parvovirus (CPV) is a small non-enveloped ssDNA virus composed of viral capsid proteins 1, 2 and 3 (VP1, VP2 and VP3). This virus has a natural affinity to cancer cells via VP2 ligands/transferrin receptors (TfRs) attachment. It has shown that VP1 and VP3 are unnecessary for capsid formation and consequently the VP2 alone is sufficient for assembly of canine parvo-virus like particle (CP-VLP) for therapeutic aims. So, in this research our purpose was to construct a recombinant bacmid shuttle vector expressing VP2 of CPV using site-specific transposition mechanism in a Bac-to-Bac baculovirus expression system. **Methods:** The mini-Tn7 transposone located in pFastBac1 donor vector containing expression cassette of CPV-VP2 was used in this experimental study that had constructed in our previous study. Firstly, the presence of gene of interest in pFastBac1 donor vector was evaluated by PCR and enzymatic digestion analysis. Then the confirmed pVP2FastBac1 plasmid was transferred into *E. coli* DH10Bac competent cells and the site-specific transposition of VP2 into a bacmid shuttle vector was accomplished using helper plasmid. Finally, the accuracy of transposition process was evaluated by a PCR panel using specific primers and PUC/M13 universal primers. **Results:** The presence of the gene of interest in pFastBac1 donor vector was confirmed by PCR and enzymatic digestion analysis and VP2-containing recombinant bacmid was subsequently constructed successfully by site-specific transposition mechanism and verified by the mentioned PCR panel. **Conclusions:** In this study, we used the Bac-to-Bac system for site-specific transposition of VP2 gene from pVP2FastBac1 to a baculovirus derived bacmid shuttle vector. The constructed recombinant bacmid can express recombinant VP2 protein in insect cells.

Received on: May 5th 2016

Revised on: May 27th 2016

Accepted on: 25th – June -2016

Published on: 10st– July-2016

KEY WORDS

Canine parvovirus, VP2, Baculovirus, Bacmid

*Corresponding author: Email: jmosavi@ihu.ac.ir; Tel.: +98 21 77104934

INTRODUCTION

The *Parvoviridae* family consists of small, icosahedral, non-enveloped viruses that contain linear single-stranded DNA (ssDNA) genomes about 5000 nucleotides long. Canine parvovirus (CPV) belongs to *Parvovirus* genus of this family and first emerged in the late 1970s as the cause of a new disease in dogs and is now prevalent in dogs worldwide [1,2,3]. CPV particles have a diameter of 25 nm and are composed of three proteins, viral protein-1 (VP1), viral protein-2 (VP2), and viral protein-3 (VP3) [4]. VP2 is the major component of the viral capsid. About 90% of the protein in the capsid is VP2, and 10% is VP1, which contains the entire VP2 sequence and 154 additional residues at its N-terminus. The third protein, VP3, is produced after intracellular proteolytic cleavage, which removes approximately 25 amino acids from the N-terminus of VP2. A wild-type capsid contains 60 subunits primarily of the VP2, along with a few VP1 and VP3 subunits [5]. CPV has a natural affinity to cancer cells via VP2 ligands/transferrin receptors (TfRs) attachment. In fact, the VP2 protein of CPV is the main part of attachment ligands for entry into specific and cancerous cells through transferrin receptors (TfRs) [6,7]. Yuan et al. (2001) claimed that VP2 can assemble into capsid-like structures and the expression of VP2 alone can result in assembly of a typically-sized virus like particle (VLP) for therapeutic aims. However, the importance of VP2 protein of canine parvovirus in binding to human cancer cells and production of veterinary detection kits for detecting the virus and also in vaccination has motivated a lot of research on production of this protein [8].

One of the best systems, considered in production of recombinant proteins, is the use of baculoviruses in insect cell expression system. The recombinant baculovirus and insect cell expression system provides high levels of recombinant proteins that undergo post-translational modifications like glycosylation. Therefore, the application of such system allows large quantity production of a desirable protein, in the native conformation [9,10]. There are various methods for construction of recombinant baculoviral vectors. One method is the use of Bac-to-Bac baculovirus expression vector system (BEVS) with an efficient site-specific transposition mechanism to generate recombinant baculovirus. This system has two major components. The pFastBac donor plasmid vector into which the gene(s) of interest will be cloned and has an expression cassette. The second component is the baculovirus shuttle vector (bacmid) into which the expression cassette will be transposed via recombinant pFastBac, constructed [11]. In this study, we tried to generate the second component of BEVS system, through construction of a recombinant bacmid DNA encoding VP2 of canine parvovirus, using site-specific transposition mechanism. This construct can be used to produce large scale of VP2 protein in insect cell.

MATERIALS AND METHODS

Plasmids, bacterial strains

The pFastBac1 (Invitrogen, USA) was used as the transfer vector. The *E. coli* strain DH5 α (Invitrogen, USA) was used for amplifying recombinant pFastBac1 donor plasmid vector. The gene of interest (VP2 gene) located in eukaryotic expression cassette of mini-Tn7 transposone of pVP2FastBac1 (generated in previous study) was available in this work. The *E. coli* strain DH10Bac (Invitrogen, USA) containing the baculovirus modified DNA (bacmid shuttle vector) with a mini-attTn7 target site and helper plasmid was used as an appropriate strain to perform the transposition process. The helper plasmid harbored by DH10Bac strains, confers resistance to tetracycline and encodes enzymes needed for transposition of the gene of interest into the bacmid.

Plasmid extraction

After the selection of proper colonies by blue/white screening, the recombinant plasmids were extracted from 1500 μ l of bacterial cell cultures using a Roche commercial kit (Germany) according to the manufacturer's instructions.

Evaluation of VP2 gene existence in pFastBac1 donor plasmid vector

The mini-Tn7 transposone located in pFastBac1 donor vector containing expression cassette of CPV-VP2 was used in this experimental study that had constructed in our previous study. So, the presence of gene of interest in expression cassette of pFastBac1 donor plasmid vector was evaluated by PCR and enzymatic digestion analysis (triple digestion using *EcoRV* and *HindIII* enzymes and also double digestion using *BamHI* and *EcoRI*) and the fragments produced were analyzed according to NEBcutter software pattern. Finally, the accuracy of the VP2 gene ORF in recombinant pFastBac1 was evaluated and confirmed by sequencing process and the analysis of sequencing results was accomplished by Chromas software, version 1.45 (data not shown).

Construction of a recombinant bacmid

The VP2-containing recombinant pFastBac1 donor plasmid was transferred into the *E. coli* DH₁₀Bac competent cells. After the transformation process, incubation for 4-6 hours was accomplished for site-specific transposition of the VP2 expression cassette from the transposing vector into the bacmid shuttle vector, leading to *lacZ* gene disruption. The presence of helper plasmid is required in this process. The transformed cells were cultured on a LB agar plate containing kanamycin (50 μ g/ml), gentamicin (7 μ g/ml), tetracycline (10 μ g/ml), X-gal (100 μ g/ml) and isopropylthio- β -galactoside (IPTG, 40 μ g/ml) and incubated at 37°C for 16h. The bacmid DNA was isolated from the overnight cultures by alkaline lysis purification method according to the general and current protocols.

It is notable that bacmid DNA is a high-molecular-weight plasmid (~ 135 kbp) and we must take care not to shear it. Over-drying, mechanically resuspension and storing the purified bacmid DNA at -20°C (as repeated freezing and thawing) may shear the DNA and gentle tapping of the bottom of the tube and storing the purified bacmid DNA at +4°C is recommended.

Analysis of recombinant bacmid DNA

The evaluation of VP2 gene existence in bacmid DNA is not performed as other plasmids. The enzymatic digestion analysis is not convenient and a PCR only by using specific primers is not sufficient. So, the transposition process accuracy and/or VP2 gene existence in bacmid DNA was evaluated by a PCR panel using both VP2 specific primers and PUC/M₁₃ universal primers (Table 1). In fact, the PCR using specific primers shows the accuracy of DH₁₀Bac transformation by recombinant pFastBac1 and the PCR using PUC/M₁₃ universal primers indicates the accuracy of recombination through site-specific transposition mechanism. The evaluation of VP2 orientation in recombinant bacmid is also indicated by the PCR using both VP2 specific and PUC/M₁₃ universal primers.

Table: 1. Details of amplified region and primer sets used for PCR analysis of recombinant bacmid

Primer Pairs	Sequence (5' to 3')	Fragment Size (bp)	The Amplified Region
pUC/M ₁₃ F VP2 specific R	5'-GTTTTCCCAGTCACGAC-3' 5'-TTAATATAATTTTCTAGGTGCTAGT-3'	3400	Tn7 R + Polyhedrin promoter + VP2 gene
VP2 specific F pUC/M ₁₃ R	5'-ATGAGTGATGGAGCAGTTCAAC-3' 5'-CAGGAAACAGCTATGAC-3'	2450	VP2 gene + Tn7 L
pUC/M ₁₃ F pUC/M ₁₃ R	5'-GTTTTCCCAGTCACGAC-3' 5'-CAGGAAACAGCTATGAC-3'	4000	Tn7 R + Polyhedrin promoter + VP2 gene + Tn7 L
VP2 specific F VP2 specific R	5'-ATGAGTGATGGAGCAGTTCAAC-3' 5'-TTAATATAATTTTCTAGGTGCTAGT-3'	1750	VP2 gene

- All the Fragment Sizes are Expressed Approximately and Has Been Calculated According to the Sequence date of the Bacmid DNA

Polymerase chain reaction programs

PCR reaction was performed in a tube containing 5 μ L of 10x PCR buffer, 1 μ L of dNTP mix (0.2 mM for each), 1.5 μ L of MgCl₂ (1.5 mM), 1 μ L of each primer with the concentration of 10 μ M or 10 pmol/ μ L (10 pmol for each), 1-2 μ L of template DNA, 1 unit of Taq DNA polymerase (Fermentas, Vilnius, Lithuania) and water nuclease-free up to 50 μ L final volume. Amplification reactions were performed in Biorad thermocycler (USA) and the PCR program included the following steps for specific primers: denaturation at 94°C for 5 min, followed by 30 cycles of 94°C for 45 sec (denaturation), 67°C for 60 sec (annealing), 72°C for 80 sec (extension), and a final extension at 72°C for 10 min. For PUC/M₁₃ universal primers the program included the following steps: denaturation at 94°C for 5 min, followed by 30 cycles of 94°C for 45 sec (denaturation), 63°C for 45 sec (annealing), 72°C for 300 sec (extension), and a final extension at 72°C for 7 min. Finally, the PCR program included the following steps for VP2 specific and PUC/M₁₃ universal primers: denaturation at 94°C for 5 min, followed by 30 cycles of 94°C for 45 sec (denaturation), 63°C for 45 sec (annealing), 72°C for 210 sec (extension), and a final extension at 72°C for 10 min. PCR products were analyzed by electrophoresis using 1% (w/v) agarose gel, stained with safe view (Kiangene, IRI).

RESULTS

Confirmation of VP2 gene existence in pFastBac1 donor plasmid vector

The presence of the gene of interest in expression cassette of pFastBac1 donor vector was confirmed by enzymatic digestion [Figures- 1(a) 1(b)] and PCR [Figure- 1b]. Triple digestion using *EcoRV* and *HindIII* enzymes and also double digestion using *BamHI* and *EcoRI* were accomplished and the fragments produced were analyzed and verified according to NEBcutter software pattern.

Analysis of recombinant bacmid DNA construct

After the verification of VP2 gene existence in expression cassette of pFastBac1, the transformation of *E. coli* DH₁₀Bac cells was accomplished successfully by VP2-containing recombinant pFastBac1 donor plasmid vector. Subsequently, the site-specific transposition of VP2 expression cassette from the transposing vector into the bacmid shuttle vector was performed with the presence of helper plasmid. After plating the cells on LB agar, the colonies containing recombinant bacmid were visible as large white colonies among the blue ones harboring the unaltered bacmids. The selected white colonies were restreaked onto a LB agar plate to ensure if they have true white phenotype. Since verification of the high molecular weight recombinant bacmid DNA is not convenient by digestion, at first a PCR using PUC/M₁₃ universal primers was performed to ensure that recombination has been done in selected colonies. All the white colonies showed a 4000 bp fragment, indicating the successful recombination in them. The results of amplification in non-recombinant bacmids of blue colonies (as the negative control) using M13/pUC primers showed a 300 bp fragment, indicating the lack of recombination performance [Figure -2].




Fig: 1. a) Enzymatic triple digestion of recombinant pFastBac1 vector (*EcoRV/HindIII*). Lane 1: GeneRuler 1 Kb DNA ladder (Fermentas, Vilnius, Lithuania). Lane 2: Expected ~ 600 bp, ~ 2500 bp and ~ 3500 bp fragments. b) PCR and Enzymatic double digestion of recombinant pFastBac1 vector (*BamHI/EcoRI*). Lane 1: VP2 gene expected fragment (1755 bp) obtained from confirmatory PCR on recombinant pFastBac1 vector. Lane 2: GeneRuler 1 Kb DNA ladder (Fermentas, Vilnius, Lithuania). Lane 3: revealed expected 1775 bp VP2 fragment and ~ 5000 bp linearized vector.




Fig: 2. Primary verification of recombination process through transposition mechanism by PCR, using PUC/M13 universal primers. Lanes 1-5: The expected fragment (~4000 bp) including Tn7 R + Polyhedrin promoter + VP2 gene + Tn7 L produced by PCR of recombinant bacmids extracted from white colonies. Lane 6: Negative control of PCR (without template). Lane 7: 1 kb DNA size marker (Yekta Tajhiz Azma, Iran). Lane 8: The expected fragment (~300 bp) produced by PCR of non-recombinant bacmid extracted from a blue colony as a negative control.

In the next step, a PCR panel was performed using VP2 specific and PUC/M₁₃ universal primers to ensure that proper transposition of the gene of interest has been done in recombinant bacmids.

The bacmid DNA contains M₁₃ forward and reverse priming sites, flanking the Tn7 mini-att site within the LacZ α -complementation region. The panel of PCR was done using PUC/M₁₃ universal forward and reverse primers, VP2 gene specific forward and reverse primers, VP2 gene specific forward primer and PUC/M₁₃ universal reverse primer and finally VP2 gene specific reverse primer and PUC/M₁₃ universal forward primer, respectively. PCR of nonrecombinant bacmid extracted from a blue colony as the negative control generated an expected ~300 bp fragment using PUC/M₁₃ universal forward and reverse primers [Figure- 3].




Fig. 3. The panel of PCR performed to confirm proper transposition of VP2 into the bacmid extracted from a white colony. Lanes 1 and 7: 1 kb DNA size marker (Yekta Tajhiz Azma, Iran). Lane 2: The expected PCR product generated using PUC/M13 universal primers (~4000 bp) including Tn7 R + Polyhedrin promoter + VP2 gene + Tn7 L fragments. Lane 3: The expected PCR product generated using PUC/M13 universal primers (~300 bp) by the non-recombinant bacmid extracted from a blue colony as a negative control. Lane 4: The PCR product generated using VP2 gene specific forward and reverse primers (1755 bp) including VP2 gene fragment. Lane 5: The PCR product generated using VP2 gene specific forward and PUC/M13 universal reverse primers (~2500 bp) including VP2 gene + Tn7 L fragments. Lane 6: The PCR product generated using PUC/M13 universal forward and VP2 gene specific reverse primers (~3500 bp) including Tn7 R + Polyhedrin promoter + VP2 gene fragments.

The results indicated the accuracy of recombination through site-specific transposition mechanism and the accuracy of VP2 orientation in the recombinant bacmid.

DISCUSSION

The expression of eukaryotic genes using baculovirus expression vectors takes advantages of their protein synthesis machinery and facilitates proper folding and post-translational modifications including glycosylation, acetylation, oligomerization and proteolysis. The Bac-to-Bac Baculovirus Expression System provides a rapid and efficient method to generate recombinant baculoviruses. This method was developed by researchers at Monsanto, and is based on site-specific transposition of an expression cassette into a baculovirus shuttle vector (bacmid) propagated in *E. coli*. The first major component of the system is a pFastBac vector into which the gene(s) of interest will be cloned. Depending on the pFastBac vector selected, expression of gene(s) of interest is controlled by the *Autographa californica* multiple nuclear polyhedrosis virus (AcMNPV) polyhedrin (PH) or p10 promoter for high-level expression in insect cells. This expression cassette is flanked by the left and right arms of Tn7, and also contains a gentamicin resistance gene and a SV40 polyadenylation signal to form a mini Tn7 [11,12]. The second major component of the System is the DH10Bac *E. coli* strain that is used as the host for pFastBac vector. DH10Bac cells contain a baculovirus shuttle vector (bacmid) with a mini-attTn7 target site and a helper plasmid. After the generation of recombinant pFastBac construct, once the pFastBac expression plasmid is transferred into DH10Bac cells, transposition occurs between the mini-Tn7 element on the pFastBac vector and the mini-attTn7 target site on the bacmid to generate a recombinant bacmid. This transposition reaction occurs in the presence of transposition proteins supplied by the helper plasmid. Once the transposition reaction is performed, we can isolate the high molecular weight recombinant bacmid DNA and transfect the bacmid DNA into insect cells to generate a recombinant baculovirus that can be used for preliminary expression experiments. After the baculoviral stock is amplified and titered, this high-titer stock can be used to infect insect cells for large-scale expression of the recombinant protein of interest [13,14].

Using the Bac-to-Bac Baculovirus Expression System to generate a recombinant baculovirus provides the following advantages over the traditional method using homologous recombination: a) Requires less than 2 weeks to identify and purify a recombinant baculovirus as compared to the 4-6 weeks required to generate a recombinant baculovirus using homologous recombination. b) Reduces the need for multiple rounds of plaque purification as the recombinant virus DNA isolated from selected colonies is not mixed with parental, non-recombinant virus. c)

Permits rapid and simultaneous isolation of multiple recombinant baculoviruses, and is suited for the expression of protein variants for structure/function studies [14].

CONCLUSION

In this study our purpose was to construct a recombinant bacmid DNA encoding viral protein-2 (VP2) of canine parvovirus using site-specific transposition mechanism. This recombinant baculoviral vector was constructed successfully under the control of polyhedrin promoter. We used the Bac-to-Bac system for site-specific transposition of VP2 gene from pVP2FastBac1 to baculovirus derived bacmid shuttle vector. The recombinant bacmid constructed here will transfect into the cultured Sf₉ (*Spodoptera frugiperda*) insect cell line to produce VP2 protein for therapeutic aims.

CONFLICT OF INTEREST

The authors declare no conflict of interests.

ACKNOWLEDGEMENT

We thank the Applied Virology Research Center in Baqiyatallah University of Medical Sciences, for financial and moral support and services provided during this part of Ph.D thesis.

FINANCIAL DISCLOSURE

None.

REFERENCES

- [1] Hueffer K, Parker JS, Weichert WS, Geisel RE, Sgro JY, Parrish CR. [2003] The natural host range shift and subsequent evolution of canine parvovirus resulted from virus-specific binding to the canine transferrin receptor, *J Virol.* 77: 1718-1726.
- [2] MacLachlan NJ, Dubovi EJ, Parvoviridae. In: MacLachlan NJ, Dubovi EJ. [2011] Editors. Fenner's Veterinary virology. 4th ed. London. Academic: 507.
- [3] Hoelzer K, Parrish CR. [2010] The emergence of parvoviruses of carnivores, *Vet Res* 41: 39–51.
- [4] Wu H, Rossmann MG. [1993] The canine parvovirus empty capsid structure, *J Mol Biol* 233(2): 231–244.
- [5] Soma T, Taharaguchi S, Ohinata T, Ishii H, Hara M. [2013] Analysis of the VP2 protein gene of canine parvovirus strains from affected dogs in Japan. *Res Vet Sci* 94: 368-371.
- [6] Dogonyaro BB, Bosman AM, Sibeko KP, Venter EH, van Vuuren M. [2013] Genetic analysis of the VP2-encoding gene of canine parvovirus strains from Africa, *Vet Microbiol* 165: 460–465.
- [7] Hueffer K, Parrish CR. [2003] Parvovirus host range, cell tropism and evolution. *Curr Opin Microbiol* 6: 392–398.
- [8] Decaro N, Desario C, Parisi A, Martella V, Lorusso A, Miccolupo A, Mari V, Colaianni ML, Cavalli A, Di Trani L, Buonavoglia C, [2009]. Genetic analysis of canine parvovirus type 2c. *Virology.* 385: 5-10.
- [9] Chang SF, Sgro JY, Parrish CR. [1992] Multiple amino acids in the capsid structure of canine parvovirus coordinately determine the canine host range and specific antigenic and hemagglutination properties. *J Virol.* 66: 6858-6867.
- [10] Wen Z, Ye L, Gao Y, Pan L, Dong K, Bu Z. [2009] Immunization by influenza virus-like particles protects aged mice against lethal influenza virus challenge. *Antiviral Res* 84(3): 215-224.
- [11] Ikonomou L, Schneider YJ, Agathos SN. [2003] Insect cell culture for industrial production of recombinant proteins. *Appl Microbiol Biotechnol* 62: 1-20.
- [12] Luckow VA, Lee SC, Barry GF, Olins PO. [1993] Efficient generation of infectious recombinant baculoviruses by site-specific transposon-mediated insertion of foreign genes into a baculovirus genome propagated in *Escherichia coli*. *J Virol* 67(8): 4566-4579.
- [13] Luckow VA, Lee SC, Barry GF, Olins PO. [1993] Efficient generation of infectious recombinant baculoviruses by site-specific transposon-mediated insertion of foreign genes into a baculovirus genome propagated in *Escherichia coli*. *J Virol.* 67(8): 4566–4579.
- [14] Sandhu KS, Naciri M, Al-Rubeai M. [2007] Prediction of recombinant protein production in an insect cell–baculovirus system using a flow cytometric technique. *J Immunol Methods.* 325(1-2): 104–113.