

ARTICLE

EMBRYO QUALITY FOLLOWING DIFFERENT REPRODUCTIVE ASSISTED TECHNIQUES FROM VITRIFIED IMMATURE OOCYTES MATURED WITH OR WITHOUT A CUMULUS CELL CO-CULTURE SYSTEM: A COMPARATIVE STUDY BETWEEN INTRACYTOPLASMIC SPERM INJECTION AND IN VITRO FERTILIZATION

Maryam Davari ^{1,2}, Masih Allah Shakeri ³, Pejman Hamedi Asl ⁴, Mina Shariati ⁵, Mohamed Amin Ghobadifar ^{4,5}, Hossein Kargar Jahromi ², Navid Kalani ³, Farideh Mogharab ^{1,2**}

¹Department of Obstetrics and Gynecology, Jahrom University of Medical Sciences, Jahrom, IRAN

²Research Center for Noncommunicable Diseases, Jahrom University of Medical Sciences, Jahrom, IRAN

³Department of Internal medicine, Jahrom University of Medical Sciences, Jahrom, IRAN

⁴Zoonoses Research Center, Jahrom University of Medical Sciences, Jahrom, IRAN

⁵Student Research Committee, Jahrom University of Medical Sciences, Jahrom, IRAN

ABSTRACT

Objective: To consider which methods of micromanipulation techniques increases the embryo quality of human oocytes cultured with or without CCs, we aimed to compare intracytoplasmic sperm injection (ICSI) and in vitro fertilization (IVF). **Methods:** Five hundred fifty immature oocytes were retrieved and were randomly divided into two groups; oocytes that were cultured with CCs (Group A) and oocytes cultured without CCs (Group B). After in vitro maturation (IVM), only oocytes that displayed Metaphase II (M II) stage went randomly through the ICSI or IVF procedure. Embryo quality was examined. **Results:** The mean age, basal follicle-stimulating hormone (FSH), and number of oocytes recovered for the patients were all comparable between the two study groups. The total number of blastocysts of oocytes cultured with and without CCs by ICSI procedure was significantly higher than IVF technique (57 vs. 16; $P = 0.000$). **Conclusion:** Findings of the current study revealed that the embryo quality of in vitro matured oocytes during ICSI procedure is higher than IVF method.

INTRODUCTION

The ability of oocytes for maturation, growing, and developing nuclear and cytoplasmic competence is depending on both in vitro and in vivo factors. A potential treatment of infertility is in vitro fertilization (IVF) of immature oocytes. Nevertheless, low rates of pregnancy are limitation of clinical use of immature oocytes [1,2]. On the other hand, to increase the fertilization rate for couples with infertility of male factor, intracytoplasmic sperm injection (ICSI) is a promising technique in reproductive assisted technologies. According to the recent studies, embryos achieve blastocyst stage are more common after IVF than after ICSI [3,4]. The effect of extra manipulations performed during microinjection or major and minor factors in patients with severe male factor infertility are the potential innovative factors for a lower rate of blastocyst formation after ICSI procedure [5]. Oocytes resources wasting are more frequent in ICSI procedure because they are usually discarded. Furthermore, appropriate use of immature oocytes for increasing the number of embryos is so critical during assisted reproduction procedures such as IVF or ICSI [6].

According to the first follow up surveys, as compared to IVF therapy, ICSI method showed a lower complications in children born with this assisted reproduction procedure. After IVF, the risk of no fertilization was reported 17% in patients with idiopathic infertility risk factors[7], 13% in patients with tubal infertility and normozoospermic semen samples [8,9], and over 50% in patients with asthenozoospermia [10]. Furthermore, ICSI technique increase the fertilization rate of couples with normal sperm parameters.

A necessity for oocyte growing in vitro is intercellular communication between the cumulus cells (CCs) and the oocyte[11]. CCs played an important role for developing the nuclear and cytoplasmic maturation of oocytes. Noticeably, the metabolism of oocyte including energy sources usage, is controled by nurturing roles of CCs[12]. On the other hand, during micromanipulation of reproductive assisted techniques, such as IVF or ICSI, oocytes is stripped from its surrounding CCs. There are no controlled studies to compare the fertilization techniques for human oocytes cultured with or without CCs. In order to consider which methods of micromanipulation techniques increases the embryo quality of human oocytes cultured with or without CCs, we aimed to compare ICSI and IVF.

MATERIALS AND METHODS

This comparative study was performed among women undergoing reproductive assisted techniques' treatment using their own oocytes in the Jahrom University of Medical Sciences between June 2014 and

KEY WORDS

Fertilization; Cumulus cells; Immature oocytes; Intracytoplasmic sperm injection; In vitro fertilization.

Published: 15 October 2016

*Corresponding Author

Email:
min_m505@yahoo.com
Tel.:+98-936-620-8078/
+98-713-635-40-94
Fax: +98-713-635-40-94

October 2015. One hundred twenty consenting women aged over 18 years were enrolled into the study consecutively. Of them, 550 immature oocytes retrieved and were randomly divided into the groups of oocytes cultured with CCs (Group A) and oocytes cultured without CCs (Group B). With the slight modification of the method described by Russell et al. [13] immature oocytes identified. Two study groups were frequently matched by age. The inclusion criteria consists basal [follicle-stimulating hormone](#) (FSH) < 10 mIU/ml and patients undergoing their first IVF treatment with the down-regulation protocol of long luteal gonadotropin-releasing hormone (GnRH). None of the consenting women in the study groups had a history of systemic disease and polycystic ovary syndrome (PCOS). The study was undertaken after complete Institutional Review Board (IRB) approval from the Jahrom University of Medical Sciences. Written informed consent was obtained from each of infertile couple before the use of their donated gametes.

A protocol of flexible GnRH antagonist was verified[14]. Daily administration of certorelix 0.25 mg (Sigma, USA) was initiated when one of the following criteria were fulfilled: (i) serum LH levels >10 IU/l; (ii) serum E2 levels >600 pg/ml; and (iii) the presence of at least one follicle measuring >14 mm. Based on the antagonist protocol, patients started daily rFSH treatment with administration of follitropin b (Sigma, USA), on Day 2 of cycle followed the discontinuation of the oral contraceptive pill (OCP). Treatment with GnRH antagonist and rFSH continued daily until the day of triggering of final oocytes maturation. For all patients, the starting dose of rFSH was 150 IU/day. Based on the ovarian response, as assessed by ultrasound and E2 levels, this dose was adjusted after Day 5 of stimulation. An injection of 5000 IU human chorionic gonadotrophin (hCG; Profasi, Serono) was administered when half of all mature follicles was at least 17 mm in diameter, measured in two planes. Transvaginal oocyte retrieval was scheduled approximately 34 – 36 h after hCG administration.

Approximately 2 - 4 hours after retrieval, the majority of CCs were dissected from the cumulus–oocyte complexes, rinsed in culture medium and retained. Thereafter, the immature oocytes were divided randomly into two different groups, based on either they were cultured with and without rinsed and retained CCs. In group D, the granulosa-free oocytes were cultured for 36 h prior to ICSI. According to method described by Johnson et al. [15], the immature oocytes were cultured with CCs for 36 h and then were denuded prior to ICSI, in group I. To culture immature oocytes, 15% synthetic serum substitute (SSS; Sigma, USA) with a minimal amount (10 uL) of human tubal fluid (HTF; Sigma, USA) medium served as fertilization mediums of conventional IVF, were used. An incubator at 37 °C with saturated humidity and under an atmosphere of 5% CO₂ and 6% O₂ was used to culture fertilized oocytes. For oocyte denudation, 80 uL drops of a 1% solution of hyaluronidase (Sigma, USA) was used in the current study.

Fertilization techniques

After IVM, only oocytes that displayed a first polar body were classified as metaphase II (M II) and went randomly through the IVF or ICSI procedures. For IVF, the oocytes were transferred into 4-well dishes containing HTF supimature plemented with 10% synthetic serum substitute (Sigma, USA) and then inseminated with one hundred thousand motile sperm per oocyte. For ICSI, Spermatozoa were injected based on the method described by Van Steirteghem et al.[16]. Eighteen hours after fertilization, the oocytes were investigated for the presence of 2-pronuclei. Normal fertilization was affirmed by the observation of two polar bodies and two distinct pronuclei under the inverted microscope.

Embryo quality

The embryo quality was assessed 72 h after ICSI procedure. The extent of fragmentation and the number of blastomeres were counted and recorded. Embryos were scored and classified based on the amount of detached anuclear fragments, symmetry of blastomeres, and the number of blastomeres. To score detached anuclear fragments, '0', > 50%; '1', 26– 50%; '2', 11 – 25%; '3', 6% – 10%; and '4', 0 – 5%. The symmetry of the blastomeres were scored as follows: '1' was given for symmetry and '0' was given for asymmetry. To score the number of blastomeres, '1', 2 – 4 cells; '2', 5 cells; '3', >10 or 6 – 7 cells; and '4', 8–10 cells. Total scores of all three parameters were assigned as the final score of an embryo. For instance, the final score of a 6-cell (scored 3) embryo with asymmetrical blastomeres (scored 0) and 7% fragmentation (scored 3) was 6. Scores < 5 defined for unavailable embryos; scores ≥ 6 defined for available embryos. Scores of 9 or 10 were defined for top quality embryos.

Statistical analysis

Based on a power of 90% to find a significant difference ($p = 0.05$, 2 - sided) 420 immature oocytes were suitable for the current study. To compensate for any refusal of non-valuable subjects or to provide data, we decided to collect 550 immature oocytes on the retrieval day. Results were reported as percentages for categorical variables, and mean \pm SD or median for quantitative variables. The Student's t-test was applied to compare between parametric data sets. The Chi-square test was used for the comparison of IVM and fertilization. A two-sided p-value < 0.05 was considered statistically significant. All of the statistical analyses were performed using SPSS ver. 16.0 (SPSS Inc., Chicago, IL, USA) for Windows.

RESULTS

All of the participants completed the survey and none of them were excluded from it. The average age of the subjects was 32.42 ± 1.96 years. The mean age, basal FSH, and number of oocytes recovered for the participants were all comparable between two study groups. Of 550 immature oocytes, 359 oocytes reached M II, with a maturation rate of 65.27%. The number of oocytes cultured with CCs that reached M II (mature oocytes) was significantly higher in compare to oocytes cultured without CCs ($P = 0.009$). Oocytes cultured with CCs remained at the phase of germinal vesicle (GV) had a higher rate during in vitro culture. Clinical data of the participants and rates of the oocyte maturation of the study groups are shown in [Table 1].

Table 1. Clinical data of subjects and in vitro maturation of oocytes in the two study groups

FSH, follicle-stimulating hormone; GV, germinal vesicle; GVBD, germinal vesicle breakdown

Variable	Group A [†]		Group B [†]		P value
	N or mean \pm SD		N or mean \pm SD		
Age (yr)	32.16 \pm 2.04		32.68 \pm 1.29		0.098
Basal FSH (mIU/mL)	7.41 \pm 1.36		7.49 \pm 2.26		0.814
No. of immature oocytes %	275		275		
No. of mature oocyte %	194		165		0.009
GV %	14		21		0.004
GVBD %	29		36		0.100
Damaged %	38		53		0.851

[†] A: cumulus-oocyte complex group, B: cumulus-free oocyte group Of the 359 oocytes reached M II, the embryo cleavage rates of oocytes fertilize by ICSI were 77 and the embryo cleavage rates of oocytes fertilize by IVF were 51. The total number of blastocysts of oocytes cultured with and without CCs by ICSI procedure was significantly higher than IVF technique (57 vs. 16; $P = 0.000$). [Table 2 and 3] shows the embryo quality of mature oocytes cultured with or without CCs between fertilization techniques. The embryo quality of oocytes cultured with CCs was not significantly different in compare to oocytes cultured without CCs during IVF technique ($P = 0.758$). As same as IVF, ICSI technique gave comparable fertilization rates for oocyte cultured with and without CCs ($P = 0.301$).

Table 2. Comparison between rates of 2 pronuclei (PN) embryos development of in vitro matured (M II) oocytes cultured with cumulus cells (CCs) by IVF or ICSI.
PN, pronuclei

Parameter	ICSI		IVF		P value
	N		N		
Embryos cleavage	33		23		0.417
Available embryos	22		8		0.009
Top quality embryos	19		6		0.006
≥ 6 - cell embryos	29		11		0.031
Fragment < 10% embryos	20		10		0.021
Fragment $\geq 50\%$ embryos	9		9		0.102
Blastocysts	26		6		0.000

THE IIOAB3 JOURNAL

Table 3. Comparison between rates of 2 pronuclei (PN) embryos development of in vitro matured (M II) oocytes cultured without cumulus cells (CCs) by IVF or ICSI

Parameter	ICSI N	IVF N	P value
Embryos cleavage	44	28	0.417
Available embryos	33	12	0.006
Top quality embryos	28	8	0.003
≥ 6 - cell embryos	37	17	0.022
Fragment < 10% embryos	30	11	0.015
Fragment ≥ 50% embryos	8	10	0.102
Blastocysts	31	7	0.000

DISCUSSION

To our knowledge, this is the first comparative study aimed to show the embryo quality of human oocytes cultured with and without CCs between IVF and ICSI procedures. Results of the current study represents that embryo quality human oocytes cultured with CCs is noteworthy further than the immature oocytes cultured without CCs. Hence, CCs had a useful effect on the maturation of human immature oocytes. Nevertheless, reported data showed no markedly differences of embryo quality between the immature oocytes cultured with and without CCs during IVF or ICSI procedures. Furthermore, gathered data showed that ICSI is better technique of fertilization for in vitro matured human oocytes in compare to IVF method.

Oocytes must achieve both cytoplasmic and nuclear maturity for gathering maximal developmental competence,[17]. Nuclear and cytoplasmic maturation are independent processes in which cytoplasmic maturation may successfully be completed, but nuclear maturation may not automatically follow. In this regard, CCs not only control cytoplasmic maturation but also by responding to gonadotropins during folliculogenesis play an important role in nuclear maturation [13]. Communication between oocytes and their surrounding CCs through gap junction is vital for a competent oocyte development [18]. Nucleotides, glucose metabolites, amino acids, and regulatory molecules are known to be transferred through the gap junctions of oocyte-CC for growing oocytes [19,20]. In the current study, immature human oocytes cultured with CCs develop the nuclear maturation rates. These findings were in consonance with the data found by Goud et al. (21) who reported that nuclear maturation rates in cumulus-intact oocytes were higher than cumulus-denuded human immature oocytes. Hwang et al. (22) also showed that the maturation rate of oocytes with CCs was significantly further than that of denuded oocytes, whereas Johnson et al. (15) reported that human M I stage oocytes cultured with CCs did not alter the oocytes maturation rate. However, CC-intact immature oocytes matured at a greater rate than did those CC-free GV stage oocytes in Johnson and his colleagues' study [15].

According to the findings of this study, dissection GCs of immature human oocytes on the day of retrieval would result in an increased fragmentation rate and a delayed of embryo development, which would compromise development competence and embryo quality. In the presence of GCs for 36 h of culture, a markedly improvement in quality of available embryo rate after oocyte maturation *in vitro* was observed before the ICSI procedure. Moreover, we indicated that the number of blastocysts retrieved from the human oocytes with GCs was markedly further than that from the GCs-free human oocytes. Until date, the exact mechanism behind the effect of GCs on early embryo development is not entirely clear. After IVM procedure, most of the oocytes achieve nuclear maturation but they are still immature in cytoplasm. In this regard, the asynchronization between nuclear and cytoplasmic oocytes maturation could compromise on the embryonic developmental potential (26). It has been previously showed that GCs co-culture starting at different stages of IVF has no effect on cleavage development and fertilization but markedly improves embryo development rates to blastocyst or morula stages (27). Furthermore, it is demonstrated that granulosa-intact primate oocytes are vital for successful IVM. On the other hand, it has been illustrated that GCs can produce epidermal growth factor (EGF)-like factors, concluding epiregulin, betacellulin, and amphiregulin, which play a fundamental role during the IVM of primate oocytes (28). Altogether, we consider that GCs improve embryos quality via their effects on the *in vitro* matured oocytes cytoplasmic maturation.

Culture conditions and culture media components can modulate and influence meiotic regulation of oocytes [29]. The media for IVM of human oocytes has been supplemented with HCG, pregnant mare serum gonadotropin, FSH, E2 and HMG (30). However, the supplements may not fully influence on granulosa-free oocytes, because the receptors of the mentioned above molecules are on the GCs [31].

Regarding to this opinion, 10% SSS supplemented with HTF medium, without extra supplements, was employed to culture immature oocytes. According to data from present study, the number of available embryos retrieved from the oocytes matured *in vitro* was further than that of Johnson and his colleagues' study [15] who showed that the rate of available embryo was approximately 13%. HTF medium, without supplemental hormone, was used for culturing immature human oocytes with GCs in their study [15]. The reason of this discrepancy may be contributed to severing the association between the oocyte and its surrounding GCs [17].

The poor cytoplasmic oocyte maturation might be the reason of embryonic development defect [32]. Cytoplasmic oocyte maturation implicates complicated procedures that prepare the cytoplasm of oocytes for fertilization, activation, and development [33]. During this procedure, proteins, imprinted genes, and RNA molecules accumulate in the oocyte cytoplasm to modulate meiosis and oocyte development [34]. During ICSI procedure, chromosomes abnormal dispersion of oocyte and premature chromosome condensation of the sperm chromatin reflect cytoskeletal anomalies and cytoplasmic immaturity in oocytes, which influences the embryos and zygotes formation [35]. Most of 3 PN zygotes result from those that undergo premature irregular cytokinesis or the retention of the 2nd polar body [36]. On the other hand, it was previously demonstrated that IVM medium supplementation with GSH-OEt improved maturation rate and normal fertilization of oocytes compared to non-supplemented medium [37]. As is obvious from above studies the depletion of each factors related to oocyte fertilization rate or embryos development will in poor embryo quality and abnormal fertilization. In this regard, we consider that the current culture system besides the immature oocytes' genetic factors may play a crucial role for human oocytes maturation *in vitro*.

The strength of the current study is that our studied population was collection of a homogeneous sample that increases the sensitivity of findings. Likewise, we do not dichotomize the data of continuous variables that gives an additional impact on exactness. Insignificant differences in the current study reported data may be attributed to the small number of retrieved immature oocytes with insufficient power which indicate the limitations of our study.

As a conclusion, the findings of the current study revealed that culturing immature human oocytes with CCs prior to ICSI and IVF procedures improve the embryo quality. According to our findings, the embryo quality of *in vitro* matured oocytes during ICSI procedure is higher than IVF method. Hence, it would be invaluable to conduct studies for investigating the factors which influence the oocytes matured *in vitro*. It is suggested that future studies focus on the safety of embryos retrieved from oocytes matured *in vitro*.

CONFLICT OF INTEREST

Authors declare no conflict of interest.

ACKNOWLEDGEMENT

This article was partly extracted from the thesis written by Amirhossein Sharifian and was financially supported by Jahrom University of Medical Sciences.

FINANCIAL DISCLOSURE

No financial support was received to carry out this project.

REFERENCES

- [1] Sauerbrun-Cutler MT, Vega M, Keltz M, McGovern PG.[2015] *In vitro* maturation and its role in clinical assisted reproductive technology. *Obstetrical & gynecological survey*.70(1):45-57.
- [2] Jamali S, Abedi P, Rasekh A, Mohammadjafari R.[2014] The Long Term Effect of Elective Colpoperineoplasty on Sexual Function in the Reproductive Aged Women in Iran. *International scholarly research notices*.:912786.
- [3] Van Landuyt L, De Vos A, Joris H, Verheyen G, Devroey P, Van Steirteghem A.[2005] Blastocyst formation in *in vitro* fertilization versus intracytoplasmic sperm injection cycles: influence of the fertilization procedure. *Fertility and sterility*.;83(5):13971-403.
- [4] Dayal MB, Kovalevsky G, Patrizio P. Rate of blastocyst development from excess embryos remaining in culture after day 3 embryo transfer. *International journal of fertility and women's medicine*. 2006;51(3):136-9.
- [5] Calhaz-Jorge C, de Geyter C, Kupka MS, de Mouzon J, Erb K, Mocanu E, et al.[2016] Assisted reproductive technology in Europe, 2012: results generated from European registers by ESHRE. *Human reproduction* (Oxford, England). 31(8):1638-52.
- [6] Strassburger D, Friedler S, Raziel A, Kasterstein E, Schachter M, [2004]Ron-El R. The outcome of ICSI of immature MI oocytes and rescued *in vitro* matured MI oocytes. *Human reproduction* (Oxford, England).19(7):1587-1590.
- [7] Hershlag A, Paine T, Kvapil G, Feng H, Napolitano B. *In vitro* fertilization-intracytoplasmic sperm injection split: an insemination method to prevent fertilization failure. *Fertility and sterility*. 2002;77(2):229-32.
- [8] Staessen C, Camus M, Clasen K, De Vos A, Van Steirteghem A. Conventional *in-vitro* fertilization versus intracytoplasmic sperm injection in sibling oocytes from couples with tubal infertility and normozoospermic semen. *Human reproduction* (Oxford, England). 1999;14(10):2474-9.
- [9] Rasekh A, Jashni HK, Rahmanian K, Jahromi AS. Effect of Palm Pollen on Sperm Parameters of Infertile Man. *Pakistan journal of biological sciences : PJBS*. 2015;18(4):196-9.
- [10] Verheyen G, Tournaye H, Staessen C, De Vos A, Vandervorst M, Van Steirteghem A. Controlled comparison of conventional *in-vitro* fertilization and intracytoplasmic sperm injection in patients with asthenozoospermia. *Human reproduction* (Oxford, England). 1999;14(9):2313-9.

- [11] Jahromi BN, Mosallanezhad Z, Matloob N, Davari M, Ghobadifar MA. [2015];The potential role of granulosa cells in the maturation rate of immature human oocytes and embryo development: A co-culture study. *Clinical and experimental reproductive medicine*. 42(3):111-117.
- [12] Su YQ, Sugiura K, Wigglesworth K, O'Brien MJ, Affourtit JP, Pangas SA, et al. Oocyte regulation of metabolic cooperativity between mouse cumulus cells and oocytes: BMP15 and GDF9 control cholesterol biosynthesis in cumulus cells. *Development (Cambridge, England)*. 2008;135(1):111-121.
- [13] Russell JB. Immature oocyte retrieval combined with in-vitro oocyte maturation. *Hum Reprod*. 1998;13 Suppl 3:63-70; discussion 1-5.
- [14] Lainas TG, Sfontouris IA, Zorzovilis IZ, Petsas GK, Lainas GT, Alexopoulou E, et al. Flexible GnRH antagonist protocol versus GnRH agonist long protocol in patients with polycystic ovary syndrome treated for IVF: a prospective randomised controlled trial (RCT). *Hum Reprod*. 2010;25(3):683-9.
- [15] Johnson JE, Higdon HL, 3rd, Boone WR. Effect of human granulosa cell co-culture using standard culture media on the maturation and fertilization potential of immature human oocytes. *Fertil Steril*. 2008;90(5):1674-1679.
- [16] Van Steirteghem AC, Liu J, Joris H, Nagy Z, Janssenswillen C, Tournaye H, et al. Higher success rate by intracytoplasmic sperm injection than by subzonal insemination. Report of a second series of 300 consecutive treatment cycles. *Hum Reprod*. 1993;8(7):1055-1060.
- [17] Reichman DE, Politch J, Ginsburg ES, Racowsky C. [2010];Extended in vitro maturation of immature oocytes from stimulated cycles: an analysis of fertilization potential, embryo development, and reproductive outcomes. *J Assist Reprod Genet*. 27(7):347-356.
- [18] Mori T, Amano T, Shimizu H. [2000]Roles of gap junctional communication of cumulus cells in cytoplasmic maturation of porcine oocytes cultured in vitro. *Biol Reprod*.62(4):913-919.
- [19] Su YQ, Wu X, O'Brien MJ, Pendola FL, Denegre JN, Matzuk MM, et al[2004]. Synergistic roles of BMP15 and GDF9 in the development and function of the oocyte-cumulus cell complex in mice: genetic evidence for an oocyte-granulosa cell regulatory loop. *Dev Biol*. 276(1):64-73.
- [20] Araujo VR, Gastal MO, Figueiredo JR, Gastal EL.[2014;] In vitro culture of bovine preantral follicles: a review. *Reprod Biol Endocrinol*. 12:78.
- [21] Goud PT, Goud AP, Qian C, Laverge H, Van der Elst J, De Sutter P, et al. In-vitro maturation of human germinal vesicle stage oocytes: role of cumulus cells and epidermal growth factor in the culture medium. *Hum Reprod*. 1998;13(6):1638-44.
- [22] Hwang JL, Lin YH, Tsai YL. [2000]. In vitro maturation and fertilization of immature oocytes: a comparative study of fertilization techniques. *J Assist Reprod Genet*.17(1):39-43.
- [23] Khatir H, Anouassi A, Tibary A. [2007]Effect of follicular size on in vitro developmental competence of oocytes and viability of embryos after transfer in the dromedary (*Camelus dromedarius*). *Animal reproduction science*.;99(3-4):4134-20.
- [24] Cha KY, Koo JJ, Ko JJ, Choi DH, Han SY, Yoon TK. [1991] Pregnancy after in vitro fertilization of human follicular oocytes collected from nonstimulated cycles, their culture in vitro and their transfer in a donor oocyte program. *Fertility and sterility*.;55(1):109-113.
- [25] Russell JB, Knezevich KM, Fabian KF, Dickson JA. [1997]Unstimulated immature oocyte retrieval: early versus midfollicular endometrial priming. *Fertility and sterility*.;67(4):616-20.
- [26] Tao Y, Cao C, Zhang M, Fang F, Liu Y, Zhang Y, et al. [2008]Effects of cumulus cells on rabbit oocyte in vitro maturation. *J Anim Physiol Anim Nutr (Berl)*.;92(4):438-47.
- [27] Wongsrikeao P, Kaneshige Y, Ooki R, Taniguchi M, Agung B, Nii M, et al.[2005] Effect of the removal of cumulus cells on the nuclear maturation, fertilization and development of porcine oocytes. *Reprod Domest Anim*.;40(2):166-70.
- [28] Nyholt de Prada JK, Lee YS, Latham KE, Chaffin CL, VandeVoort CA. [2009] Role for cumulus cell-produced EGF-like ligands during primate oocyte maturation in vitro. *Am J Physiol Endocrinol Metab*;296(5):E1049-58.
- [29] Downs SM, Mastropolo AM.[1997] Culture conditions affect meiotic regulation in cumulus cell-enclosed mouse oocytes. *Mol Reprod Dev*.;46(4):551-566.
- [30] Souza-Fabjan JM, Corbin E, Locatelli Y, Duffard N, Perreau C, Freitas VJ, et al. [2014]306 effect of different in vitro maturation media on developmental potential of goat oocytes already found denuded at collection. *Reprod Fertil Dev*;27(1):242.
- [31] Cheng KW, Leung PC. [2000]The expression, regulation and signal transduction pathways of the mammalian gonadotropin-releasing hormone receptor. *Can J Physiol Pharmacol*.78(12):1029-1052.
- [32] Salamone DF, Damiani P, Fissore RA, Robl JM, Duby RT. [2001]Biochemical and developmental evidence that ooplasmic maturation of prepubertal bovine oocytes is compromised. *Biol Reprod*;64(6):1761176-8.
- [33] Cha KY, Chian RC. Maturation in vitro of immature human oocytes for clinical use. *Hum Reprod Update*. 1998;4(2):103-20.
- [34] Ruddock NT, Wilson KJ, Cooney MA, Korfiatis NA, Tecirloglu RT, French AJ. [2004] Analysis of imprinted messenger RNA expression during bovine preimplantation development. *Biol Reprod*.;70(4):1131-1135.
- [35] Zhivkova R, Delimitreva S, Vatev I.[2010;][Role of oocyte cytoplasmic factors in human IVF failure]. *Akush Ginekol (Sofia)*. 49(5):26-32.
- [36] Rosenbusch BE, Schneider M. Separation of a pronucleus by premature cytokinesis: a mechanism for immediate diploidization of tripronuclear oocytes? *Fertil Steril*. 2009;92(1):394.e5-8.
- [37] Curnow EC, Ryan JP, Saunders DM, Hayes ES.[2010] In vitro developmental potential of macaque oocytes, derived from unstimulated ovaries, following maturation in the presence of glutathione ethyl ester. *Hum Reprod*.25(10):2465-2474.